



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Development of banana pollen.—An extensive investigation²¹ of three races of the edible banana (*Musa sapientum*) has shown that they can be distinguished by the number of chromosomes, "*Dole*" having 8, "*Radjah Siam*" 16, and "*Kladi*" 24, as the haploid numbers, so that the races might be designated as vars. *univalens*, *bivalens*, and *trivalens*. The volume of the nuclei, but not their surfaces, is in the ratio 1:2:3. With the increase in the number of chromosomes came disturbances in the development of pollen, some of the chromosomes not passing to the poles, but remaining behind and forming extra nuclei. The size of the tetrad varies in a given anther, although the number of chromosomes in the entire tetrad is constant. Sometimes as many as eight pollen grains are formed from a single mother cell.

Prochromosomes are easily distinguished in the pollen mother cell, and in *Musa Dole* TISCHLER was able to show that the number of prochromosomes was equal to the diploid number of chromosomes. Probably there is a fusion of prochromosomes at synapsis. The splitting of chromosomes at the streptonema stage TISCHLER regards as genuine and not merely apparent.—CHARLES J. CHAMBERLAIN.

Parthenogenesis in Taraxacum.—Parthenogenesis in *Taraxacum* has been investigated again, this time by SCHKORBATOW²² who writes in Russian, but adds a summary in German, from which the following points are taken: The removal of anthers does not in any way affect the germination of seeds. Various colors of seeds, like clear green and dark brown, may become fixed and hereditary. At metaphase of the first division in the embryo sac, the chromosomes show various and characteristic forms, but the chromosomes seldom take the arrangement belonging to the heterotypic mitosis, and when they do, the author regards the phenomena as atavistic. Amitotic divisions occur in the embryo sac, in the endosperm, and in early stages of the embryo, in the last case all the nuclei but one becoming resorbed, so that the cells are left uninucleate.—CHARLES J. CHAMBERLAIN.

The origin of the vacuole.—Probably most botanists believe that the large vacuoles of plants arise by the coalescence of numerous smaller ones. A paper by BENSLEY,²³ dealing with the canalicular apparatus of animals, gives also a description of root tips and the tapetum of anthers. The fixing agent used was: neutral formalin (freshly distilled), 10 cc.; water, 90 cc.; potassium bichromate, 2.5 g.; mercuric chloride, 5.0 g. With this fixing

²¹ TISCHLER, G., Untersuchungen über die Entwicklung des Bananen-Pollens. I. Archiv. für Zellforschung 5:622-670. pls. 30, 31. 1910.

²² SCHKORBATOW, L., Parthenogenetische und apogame Entwicklung bei den Blütenpflanzen Entwicklungsgeschichtliche Studien an *Taraxacum officinale* Wigg. Bot. Institut Charkow. pp. 43. pl. 1. figs. 4. 1910.

²³ BENSLEY, R. R., On the nature of the canalicular apparatus of animal cells. Biol. Bull. 19:174-194. figs. 1-3. 1910.